

Eastern Illinois University
The Keep

Faculty Research and Creative Activity

Chemistry

January 2010

10,12-Dimethylpteridino[6,7-f][1,10]-phenanthroline-11,13(10H,12H)-dione–chloroform (1/1)

Waynie Olapratath
Eastern Illinois University

Jennifer Rodin
Eastern Illinois University

Kraig A. Wheeler
Eastern Illinois University, kawheeler@eiu.edu

Mark E. McGuire
Eastern Illinois University

Follow this and additional works at: http://thekeep.eiu.edu/chemistry_fac

 Part of the [Organic Chemistry Commons](#)

Recommended Citation

Olapratath, Waynie; Rodin, Jennifer; Wheeler, Kraig A.; and McGuire, Mark E., "10,12-Dimethylpteridino[6,7-f][1,10]-phenanthroline-11,13(10H,12H)-dione–chloroform (1/1)" (2010). *Faculty Research and Creative Activity*. 2.
http://thekeep.eiu.edu/chemistry_fac/2

This Article is brought to you for free and open access by the Chemistry at The Keep. It has been accepted for inclusion in Faculty Research and Creative Activity by an authorized administrator of The Keep. For more information, please contact tabrunns@eiu.edu.

10,12-Dimethylpteridino[6,7-f][1,10]-phenanthroline-11,13(10H,12H)-dione-chloroform (1/1)

Wayne Olaprat, Jennifer Roden, Kraig A. Wheeler and Mark E. McGuire*

Department of Chemistry, Eastern Illinois University, 600 Lincoln Ave., Charleston, IL 61920, USA

Correspondence e-mail: memmcguire@eiu.edu


Received 22 July 2010; accepted 5 August 2010

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.032; wR factor = 0.095; data-to-parameter ratio = 12.3.

In the title co-crystal, $\text{C}_{18}\text{H}_{12}\text{N}_6\text{O}_2\cdot\text{CHCl}_3$, intramolecular $\text{Cl}_3\text{C}-\text{H}\cdots\text{N}$ hydrogen-bonding interactions occur between a single CHCl_3 and both N atoms at the 1,10-positions on the phenanthroline portion of the molecule. The interplanar distance between inversion-related molecules is $3.241(2)\text{ \AA}$.

Related literature

For the synthesis, see: Black *et al.* (1993). For the possible use of metal complexes of this ligand as DNA probes, see: Gao *et al.* (2007); Lawrence *et al.* (2006). For studies involving the non-methylated analog of the title compound, see: Chen *et al.* (2010); Dalton *et al.* (2008); Ozawa *et al.* (2006). For a related structure, see: Ton & Bolte (2005). For $\text{Cl}_3\text{C}-\text{H}\cdots\text{N}$ hydrogen bonding, see: Fan *et al.* (2009); Li & Wang (2007).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{12}\text{N}_6\text{O}_2\cdot\text{CHCl}_3$
 $M_r = 463.70$
 Monoclinic, $P2_1/n$

$a = 8.9043(2)\text{ \AA}$
 $b = 16.4009(4)\text{ \AA}$
 $c = 13.4872(4)\text{ \AA}$

$\beta = 108.058(1)^\circ$
 $V = 1872.63(8)\text{ \AA}^3$
 $Z = 4$
 Cu $K\alpha$ radiation

$\mu = 4.72\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.45 \times 0.22 \times 0.17\text{ mm}$

Data collection

Bruker APEXII CCD
 diffractometer
 Absorption correction: numerical
 (*SADABS*; Bruker, 2008)
 $T_{\min} = 0.224$, $T_{\max} = 0.508$

15543 measured reflections
 3370 independent reflections
 3115 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.095$
 $S = 1.08$
 3370 reflections

273 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.37\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.26\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C19—H19···N1	1.00	2.39	3.188 (2)	136
C19—H19···N6	1.00	2.26	3.181 (2)	152

Data collection: *APEX2* (Bruker, 2008); cell refinement: *APEX2* and *SAINT* (Bruker, 2008); data reduction: *SAINT* and *XPREP* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *X-SEED*.

We gratefully acknowledge the financial support of the National Science Foundation (CHE-0722547) to KAW.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FL2311).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Black, K. J., Huang, H., High, S., Starks, L., Olson, M. & McGuire, M. E. (1993). *Inorg. Chem.* **32**, 5591–5596.
- Bruker (2008). *APEX2*, *SADABS*, *SAINT* and *XPREP*. Bruker AXS, Inc., Madison, Wisconsin, USA.
- Chen, X., Gao, F., Zhou, Z.-X., Yang, W.-Y., Guo, L.-T. & Ji, L.-N. (2010). *J. Inorg. Biochem.* **104**, 576–582.
- Dalton, S. R., Glazier, S., Leung, B., Win, S., Megatulski, C. & Nieter Burgmayer, S. J. (2008). *J. Biol. Inorg. Chem.* **13**, 1133–1148.
- Fan, H., Molivia, A. C. D., Eliason, J. K., Olson, J. L., Green, D. D., Gealy, M. W. & Ulness, D. J. (2009). *Chem. Phys. Lett.* **479**, 43–46.
- Gao, F., Chao, H., Zhou, F., Xu, L.-C., Zheng, K.-C. & Ji, L.-N. (2007). *Helv. Chim. Acta* **90**, 36–50.
- Lawrence, D., Vaidyanathan, V. G. & Nair, B.-U. (2006). *J. Inorg. Biochem.* **100**, 1244–1251.
- Li, A. Y. & Wang, S. W. (2007). *J. Mol. Struct.* **807**, 191–199.
- Ozawa, T., Kishi, Y., Miyamoto, K., Wasada-Tsutsui, Y., Funahashi, Y., Jitsukawa, K. & Masuda, H. (2006). *Adv. Mater. Res.* **11–12**, 277–280.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Ton, Q. C. & Bolte, M. (2005). *Acta Cryst. E* **61**, o1406–o1407.

supplementary materials

Acta Cryst. (2010). E66, o2288 [doi:10.1107/S1600536810031570]

10,12-Dimethylpteridino[6,7-f][1,10]phenanthroline-11,13(10H,12H)-dione-chloroform (1/1)

W. Olapratth, J. Roden, K. A. Wheeler and M. E. McGuire

Comment

We first reported the synthesis of the title compound as part of our effort to study the pH-dependent electron transfer reactions of transition metal complexes of riboflavin mimics (Black *et al.*, 1993). Since that time, others have investigated metal complexes of this ligand as possible DNA probes (see, for example, Gao *et al.*, 2007; Lawrence *et al.*, 2006). The title compound is only slightly soluble in many common solvents; it appears at least moderately soluble in CHCl₃. We have grown crystals by vapor diffusion of hexane into a CHCl₃/95% EtOH solution. As shown in Figure 1, the crystal structure revealed interesting H-bonding interactions involving the H-atom on CHCl₃ and the N-atoms in the 1,10-phenanthroline portion of the ligand (C19—H···N1: 3.188 (2) Å, 136° and C19—H···N6: 3.181 (2) Å, 152°). This type of interaction has been reported previously (Ton & Bolte, 2005) for a co-crystal of CHCl₃ and 1,10-phenanthroline(Cl₃C—H···N: 3.175 (3) Å, 146° and 3.225 (3) Å, 141°). *Ab initio* calculations on the interaction between CHCl₃ and pyridine (Li & Wang, 2007) predict a range of Cl₃C—H···N distances (3.13–3.25 Å) and angles (156–167°), depending on the basis sets used. Fan *et al.* (2009) reported very weak Cl₃C—H···N interaction in pyridine/chloroform solutions as measured by anti-Stokes Raman scattering. The interplanar distance between inversion-related molecules in **I** (viewed along the *a*-axis) is 3.241 (2) Å. The molecular planes in alternating stacks are oriented at 12.46 (2)° to each other; this could at least partially be due to packing constraints imposed by the slightly out-of-plane C18-methyl group (see packing diagram in Figure 2). Studies involving the non-methylated analog of the title compound have also been reported (Chen *et al.*, 2010; Dalton *et al.*, 2008; Gao *et al.*, 2007; Ozawa *et al.*, 2006) although, to our knowledge, no crystal structures of this analog have been reported.

Experimental

The title compound was prepared and purified using a previously published procedure (Black *et al.*, 1993). A small amount of this compound was dissolved in ~1 ml of CHCl₃ and one drop of 95% ethanol was added to the mixture. The vial containing this mixture was placed in a beaker with ~1 ml of hexane and the beaker was loosely sealed. In 14 days, yellow crystals, suitable for data collection, were observed on the sides of the vial.

Refinement

H atoms were positioned geometrically with C—H = 0.95, 0.96 and 1.00 Å, for aromatic, methyl and methine H atoms, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2\text{U}_{\text{eq}}(\text{C})$ [$U_{\text{iso}}(\text{H}) = 1.5\text{U}_{\text{eq}}(\text{C})$ for methyl groups].

supplementary materials

Figures




Fig. 1. Perspective view of the title compound, with the atom numbering; displacement ellipsoids are at the 50% probability level.




Fig. 2. Packing diagram for the title compound. The molecular planes of alternating stacks of molecules are oriented at 12.46 (2) $^{\circ}$ relative to one another. The CHCl₃ molecules and H-atoms have been omitted for clarity.

10,12-Dimethylpteridino[6,7-f][1,10]phenanthroline-11,13(10H,\ 12H)-dione–chloroform (1/1)

Crystal data

C₁₈H₁₂N₆O₂·CHCl₃

F(000) = 944

M_r = 463.70

D_x = 1.645 Mg m⁻³

Monoclinic, *P*2₁/*n*

Cu *K* α radiation, λ = 1.54178 Å

Hall symbol: -P 2yn

Cell parameters from 8311 reflections

a = 8.9043 (2) Å

θ = 4.3–71.9 $^{\circ}$

b = 16.4009 (4) Å

μ = 4.72 mm⁻¹

c = 13.4872 (4) Å

T = 173 K

β = 108.058 (1) $^{\circ}$

Transparent prism, yellow

V = 1872.63 (8) Å³

0.45 × 0.22 × 0.17 mm

Z = 4

Data collection

Bruker APEXII CCD
diffractometer

3370 independent reflections

Radiation source: fine-focus sealed tube
graphite

3115 reflections with $I > 2\sigma(I)$

phi and ω scans

R_{int} = 0.033

Absorption correction: numerical
(SADABS; Bruker, 2008)

θ_{\max} = 68.2 $^{\circ}$, θ_{\min} = 4.4 $^{\circ}$

T_{\min} = 0.224, T_{\max} = 0.508

h = -10→10

15543 measured reflections

k = -19→19

l = -16→15

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct
methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)]$ = 0.032

Hydrogen site location: inferred from neighbouring
sites

$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.061P)^2 + 0.6946P]$
3370 reflections	where $P = (F_o^2 + 2F_c^2)/3$
273 parameters	$(\Delta/\sigma)_{\max} = 0.001$
0 restraints	$\Delta\rho_{\max} = 0.37 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	1.11933 (6)	0.91490 (3)	0.06882 (4)	0.03314 (15)
Cl2	1.02227 (5)	0.75947 (3)	0.13113 (3)	0.02516 (14)
Cl3	0.78834 (5)	0.87388 (3)	0.00893 (4)	0.03132 (15)
O1	0.58359 (16)	0.89641 (8)	0.76117 (10)	0.0252 (3)
O2	0.50035 (16)	1.16797 (8)	0.78817 (10)	0.0253 (3)
N1	0.93260 (18)	0.83887 (9)	0.33300 (11)	0.0211 (3)
N2	0.69578 (17)	0.92402 (9)	0.59325 (11)	0.0185 (3)
N3	0.53018 (17)	1.03163 (9)	0.76877 (11)	0.0199 (3)
N4	0.61174 (18)	1.12873 (9)	0.66532 (11)	0.0193 (3)
N5	0.71061 (17)	1.08699 (9)	0.53254 (11)	0.0185 (3)
N6	0.94027 (18)	0.99643 (9)	0.27110 (11)	0.0206 (3)
C1	0.9330 (2)	0.76237 (11)	0.36516 (14)	0.0228 (4)
H1	0.9744	0.7215	0.3310	0.027*
C2	0.8759 (2)	0.73847 (11)	0.44641 (15)	0.0232 (4)
H2	0.8798	0.6829	0.4670	0.028*
C3	0.8141 (2)	0.79660 (11)	0.49591 (14)	0.0207 (4)
H3	0.7739	0.7820	0.5510	0.025*
C4	0.8113 (2)	0.87791 (10)	0.46354 (13)	0.0180 (3)
C5	0.7528 (2)	0.94334 (10)	0.51497 (13)	0.0181 (3)
C6	0.6493 (2)	0.98516 (11)	0.64097 (13)	0.0187 (4)
C7	0.5877 (2)	0.96493 (10)	0.72797 (13)	0.0193 (4)
C8	0.5445 (2)	1.11330 (11)	0.74324 (13)	0.0202 (4)
C9	0.6584 (2)	1.06698 (10)	0.61137 (13)	0.0185 (4)
C10	0.7584 (2)	1.02511 (10)	0.48401 (13)	0.0179 (3)
C11	0.8194 (2)	1.04465 (11)	0.39814 (13)	0.0186 (3)

supplementary materials

C12	0.8254 (2)	1.12494 (11)	0.36400 (14)	0.0211 (4)
H12	0.7857	1.1686	0.3949	0.025*
C13	0.8895 (2)	1.13969 (11)	0.28512 (14)	0.0221 (4)
H13	0.8958	1.1936	0.2609	0.027*
C14	0.9452 (2)	1.07352 (11)	0.24152 (14)	0.0226 (4)
H14	0.9895	1.0843	0.1872	0.027*
C15	0.8776 (2)	0.98147 (10)	0.34867 (13)	0.0185 (3)
C16	0.8732 (2)	0.89648 (11)	0.38207 (13)	0.0181 (4)
C17	0.4656 (2)	1.01675 (11)	0.85485 (14)	0.0241 (4)
H17A	0.4159	0.9628	0.8465	0.036*
H17B	0.3867	1.0586	0.8543	0.036*
H17C	0.5511	1.0189	0.9213	0.036*
C18	0.6292 (2)	1.21489 (11)	0.64044 (14)	0.0232 (4)
H18A	0.7078	1.2195	0.6034	0.035*
H18B	0.6641	1.2466	0.7051	0.035*
H18C	0.5275	1.2359	0.5962	0.035*
C19	0.9722 (2)	0.86285 (11)	0.10784 (14)	0.0226 (4)
H19	0.9645	0.8881	0.1736	0.027*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0291 (3)	0.0310 (3)	0.0357 (3)	-0.00401 (19)	0.0049 (2)	0.01122 (19)
Cl2	0.0350 (3)	0.0176 (2)	0.0227 (2)	0.00243 (17)	0.00862 (19)	0.00046 (15)
Cl3	0.0251 (3)	0.0264 (3)	0.0357 (3)	0.00431 (18)	-0.0004 (2)	-0.00619 (18)
O1	0.0364 (7)	0.0162 (6)	0.0245 (7)	-0.0019 (5)	0.0115 (6)	-0.0004 (5)
O2	0.0348 (7)	0.0187 (6)	0.0233 (6)	0.0043 (5)	0.0104 (6)	-0.0023 (5)
N1	0.0255 (8)	0.0172 (7)	0.0193 (7)	-0.0007 (6)	0.0048 (6)	-0.0020 (6)
N2	0.0213 (7)	0.0153 (7)	0.0167 (7)	-0.0020 (6)	0.0023 (6)	-0.0017 (5)
N3	0.0240 (7)	0.0175 (7)	0.0179 (7)	-0.0003 (6)	0.0059 (6)	-0.0019 (5)
N4	0.0258 (8)	0.0133 (7)	0.0179 (7)	0.0018 (6)	0.0054 (6)	-0.0010 (5)
N5	0.0213 (7)	0.0146 (7)	0.0170 (7)	0.0009 (5)	0.0023 (6)	0.0001 (5)
N6	0.0255 (8)	0.0180 (7)	0.0168 (7)	-0.0011 (6)	0.0042 (6)	0.0004 (6)
C1	0.0286 (9)	0.0162 (9)	0.0228 (9)	0.0007 (7)	0.0067 (8)	-0.0030 (7)
C2	0.0296 (9)	0.0141 (8)	0.0244 (9)	0.0000 (7)	0.0065 (8)	0.0000 (7)
C3	0.0239 (9)	0.0179 (9)	0.0189 (8)	-0.0021 (7)	0.0047 (7)	-0.0011 (7)
C4	0.0194 (8)	0.0150 (8)	0.0165 (8)	-0.0018 (6)	0.0011 (7)	-0.0015 (6)
C5	0.0194 (8)	0.0161 (8)	0.0159 (8)	-0.0021 (7)	0.0015 (7)	-0.0014 (6)
C6	0.0201 (8)	0.0168 (8)	0.0167 (8)	-0.0007 (7)	0.0019 (7)	-0.0014 (6)
C7	0.0227 (8)	0.0164 (9)	0.0164 (8)	-0.0023 (7)	0.0025 (7)	-0.0026 (6)
C8	0.0221 (9)	0.0183 (8)	0.0172 (8)	0.0007 (7)	0.0018 (7)	-0.0006 (7)
C9	0.0193 (8)	0.0164 (8)	0.0165 (8)	-0.0002 (7)	0.0006 (7)	-0.0021 (6)
C10	0.0184 (8)	0.0161 (8)	0.0163 (8)	-0.0001 (6)	0.0011 (7)	-0.0017 (6)
C11	0.0198 (8)	0.0168 (8)	0.0161 (8)	-0.0002 (7)	0.0012 (7)	0.0006 (6)
C12	0.0228 (9)	0.0172 (9)	0.0198 (9)	0.0009 (7)	0.0015 (7)	-0.0010 (7)
C13	0.0256 (9)	0.0175 (8)	0.0201 (9)	-0.0015 (7)	0.0024 (7)	0.0033 (7)
C14	0.0266 (9)	0.0209 (9)	0.0187 (8)	-0.0027 (7)	0.0047 (7)	0.0017 (7)
C15	0.0206 (8)	0.0163 (9)	0.0154 (8)	-0.0007 (7)	0.0008 (7)	-0.0003 (6)

C16	0.0193 (8)	0.0158 (8)	0.0156 (8)	-0.0006 (7)	0.0000 (7)	-0.0018 (6)
C17	0.0314 (10)	0.0217 (9)	0.0210 (9)	-0.0015 (7)	0.0105 (8)	-0.0015 (7)
C18	0.0339 (10)	0.0139 (8)	0.0218 (9)	0.0014 (7)	0.0084 (8)	-0.0002 (7)
C19	0.0264 (9)	0.0191 (8)	0.0197 (9)	-0.0006 (7)	0.0036 (7)	-0.0005 (7)

Geometric parameters (\AA , $^\circ$)

C1—C19	1.7747 (18)	C3—C4	1.401 (2)
Cl2—C19	1.7564 (18)	C3—H3	0.9500
Cl3—C19	1.7716 (19)	C4—C16	1.407 (2)
O1—C7	1.214 (2)	C4—C5	1.459 (2)
O2—C8	1.214 (2)	C5—C10	1.410 (2)
N1—C1	1.327 (2)	C6—C9	1.410 (2)
N1—C16	1.352 (2)	C6—C7	1.479 (2)
N2—C6	1.325 (2)	C10—C11	1.459 (2)
N2—C5	1.343 (2)	C11—C12	1.401 (2)
N3—C7	1.392 (2)	C11—C15	1.415 (2)
N3—C8	1.399 (2)	C12—C13	1.376 (3)
N3—C17	1.467 (2)	C12—H12	0.9500
N4—C9	1.384 (2)	C13—C14	1.396 (3)
N4—C8	1.384 (2)	C13—H13	0.9500
N4—C18	1.472 (2)	C14—H14	0.9500
N5—C9	1.327 (2)	C15—C16	1.469 (2)
N5—C10	1.346 (2)	C17—H17A	0.9800
N6—C14	1.331 (2)	C17—H17B	0.9800
N6—C15	1.352 (2)	C17—H17C	0.9800
C1—C2	1.399 (3)	C18—H18A	0.9800
C1—H1	0.9500	C18—H18B	0.9800
C2—C3	1.373 (3)	C18—H18C	0.9800
C2—H2	0.9500	C19—H19	1.0000
C1—N1—C16	117.55 (15)	N5—C10—C11	118.16 (15)
C6—N2—C5	117.01 (15)	C5—C10—C11	120.02 (15)
C7—N3—C8	125.62 (15)	C12—C11—C15	118.44 (16)
C7—N3—C17	117.70 (15)	C12—C11—C10	121.93 (16)
C8—N3—C17	116.25 (14)	C15—C11—C10	119.62 (15)
C9—N4—C8	122.44 (15)	C13—C12—C11	119.12 (16)
C9—N4—C18	120.82 (14)	C13—C12—H12	120.4
C8—N4—C18	116.73 (14)	C11—C12—H12	120.4
C9—N5—C10	116.45 (15)	C12—C13—C14	118.39 (16)
C14—N6—C15	117.66 (15)	C12—C13—H13	120.8
N1—C1—C2	123.77 (16)	C14—C13—H13	120.8
N1—C1—H1	118.1	N6—C14—C13	124.28 (16)
C2—C1—H1	118.1	N6—C14—H14	117.9
C3—C2—C1	118.95 (17)	C13—C14—H14	117.9
C3—C2—H2	120.5	N6—C15—C11	122.09 (16)
C1—C2—H2	120.5	N6—C15—C16	117.76 (15)
C2—C3—C4	118.70 (16)	C11—C15—C16	120.14 (15)
C2—C3—H3	120.7	N1—C16—C4	122.58 (16)
C4—C3—H3	120.7	N1—C16—C15	117.50 (15)

supplementary materials

C3—C4—C16	118.44 (16)	C4—C16—C15	119.91 (15)
C3—C4—C5	121.75 (15)	N3—C17—H17A	109.5
C16—C4—C5	119.76 (16)	N3—C17—H17B	109.5
N2—C5—C10	120.93 (16)	H17A—C17—H17B	109.5
N2—C5—C4	118.54 (15)	N3—C17—H17C	109.5
C10—C5—C4	120.52 (15)	H17A—C17—H17C	109.5
N2—C6—C9	121.92 (16)	H17B—C17—H17C	109.5
N2—C6—C7	117.64 (16)	N4—C18—H18A	109.5
C9—C6—C7	120.43 (16)	N4—C18—H18B	109.5
O1—C7—N3	121.63 (16)	H18A—C18—H18B	109.5
O1—C7—C6	124.15 (16)	N4—C18—H18C	109.5
N3—C7—C6	114.20 (15)	H18A—C18—H18C	109.5
O2—C8—N4	121.82 (16)	H18B—C18—H18C	109.5
O2—C8—N3	121.06 (16)	Cl2—C19—Cl3	110.94 (10)
N4—C8—N3	117.12 (15)	Cl2—C19—Cl1	110.33 (10)
N5—C9—N4	118.51 (16)	Cl3—C19—Cl1	108.81 (10)
N5—C9—C6	121.85 (16)	Cl2—C19—H19	108.9
N4—C9—C6	119.65 (16)	Cl3—C19—H19	108.9
N5—C10—C5	121.81 (16)	Cl1—C19—H19	108.9
C16—N1—C1—C2	-0.1 (3)	N2—C6—C9—N5	1.6 (3)
N1—C1—C2—C3	0.7 (3)	C7—C6—C9—N5	-178.85 (16)
C1—C2—C3—C4	-0.4 (3)	N2—C6—C9—N4	-178.51 (16)
C2—C3—C4—C16	-0.5 (3)	C7—C6—C9—N4	1.0 (3)
C2—C3—C4—C5	-177.73 (17)	C9—N5—C10—C5	0.3 (2)
C6—N2—C5—C10	-1.6 (2)	C9—N5—C10—C11	-178.90 (15)
C6—N2—C5—C4	177.43 (15)	N2—C5—C10—N5	1.4 (3)
C3—C4—C5—N2	-2.1 (3)	C4—C5—C10—N5	-177.55 (16)
C16—C4—C5—N2	-179.32 (15)	N2—C5—C10—C11	-179.40 (15)
C3—C4—C5—C10	176.93 (16)	C4—C5—C10—C11	1.6 (3)
C16—C4—C5—C10	-0.3 (3)	N5—C10—C11—C12	-1.6 (3)
C5—N2—C6—C9	0.1 (3)	C5—C10—C11—C12	179.22 (16)
C5—N2—C6—C7	-179.42 (15)	N5—C10—C11—C15	177.03 (16)
C8—N3—C7—O1	173.01 (17)	C5—C10—C11—C15	-2.2 (2)
C17—N3—C7—O1	0.8 (3)	C15—C11—C12—C13	-1.0 (3)
C8—N3—C7—C6	-8.3 (2)	C10—C11—C12—C13	177.63 (16)
C17—N3—C7—C6	179.50 (15)	C11—C12—C13—C14	0.5 (3)
N2—C6—C7—O1	3.3 (3)	C15—N6—C14—C13	-0.2 (3)
C9—C6—C7—O1	-176.22 (17)	C12—C13—C14—N6	0.1 (3)
N2—C6—C7—N3	-175.31 (15)	C14—N6—C15—C11	-0.3 (3)
C9—C6—C7—N3	5.1 (2)	C14—N6—C15—C16	-179.49 (15)
C9—N4—C8—O2	-178.08 (16)	C12—C11—C15—N6	1.0 (3)
C18—N4—C8—O2	0.8 (3)	C10—C11—C15—N6	-177.72 (16)
C9—N4—C8—N3	2.3 (3)	C12—C11—C15—C16	-179.93 (16)
C18—N4—C8—N3	-178.82 (15)	C10—C11—C15—C16	1.4 (2)
C7—N3—C8—O2	-174.77 (17)	C1—N1—C16—C4	-0.8 (3)
C17—N3—C8—O2	-2.5 (2)	C1—N1—C16—C15	178.08 (16)
C7—N3—C8—N4	4.9 (3)	C3—C4—C16—N1	1.1 (3)
C17—N3—C8—N4	177.20 (15)	C5—C4—C16—N1	178.42 (15)
C10—N5—C9—N4	178.39 (15)	C3—C4—C16—C15	-177.78 (16)

C10—N5—C9—C6	−1.7 (2)	C5—C4—C16—C15	−0.5 (3)
C8—N4—C9—N5	174.92 (15)	N6—C15—C16—N1	0.1 (2)
C18—N4—C9—N5	−4.0 (2)	C11—C15—C16—N1	−179.05 (15)
C8—N4—C9—C6	−5.0 (3)	N6—C15—C16—C4	179.05 (15)
C18—N4—C9—C6	176.14 (16)	C11—C15—C16—C4	−0.1 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C19—H19···N1	1.00	2.39	3.188 (2)	136
C19—H19···N6	1.00	2.26	3.181 (2)	152

supplementary materials

Fig. 1




Fig. 2

