

November 2006

(Acetonitrile- κ N){2-[bis(2-pyridylethyl)-
amino]ethanol- κ 4N,N',N'',O}zinc(II) bis-
(perchlorate) monohydrate

Hong-Chang Liang
San Diego State University

Marcel M. Hetu
San Diego State University

Kraig A. Wheeler
Eastern Illinois University, kawheeler@eiu.edu

Lev N. Zakharov
University of California - San Diego

Arnold L. Rheingold
University of California - San Diego

Follow this and additional works at: http://thekeep.eiu.edu/chemistry_fac

 Part of the [Organic Chemistry Commons](#)

Recommended Citation

Liang, Hong-Chang; Hetu, Marcel M.; Wheeler, Kraig A.; Zakharov, Lev N.; and Rheingold, Arnold L., "(Acetonitrile- κ N){2-[bis(2-pyridylethyl)amino]ethanol- κ 4N,N',N'',O}zinc(II) bis(perchlorate) monohydrate" (2006). *Faculty Research and Creative Activity*. 13.

http://thekeep.eiu.edu/chemistry_fac/13

(Acetonitrile- κN)[2-[bis(2-pyridyl-ethyl)amino]ethanol- $\kappa^4 N, N', N'', O$]-zinc(II) bis(perchlorate) monohydrateHong-Chang Liang,^{a*} Marcel M. Hetu,^a Kraig A. Wheeler,^b Lev N. Zakharov^c and Arnold L. Rheingold^c

^aDepartment of Chemistry and Biochemistry, San Diego State University, 5500 Campanile Drive, San Diego, CA 92182-1030, USA, ^bDepartment of Chemistry, Eastern Illinois University, 600 Lincoln Avenue, Charleston, IL 61920-3099, USA, and ^cDepartment of Chemistry and Biochemistry, University of California at San Diego, La Jolla, CA 92093-0358, USA
Correspondence e-mail: hcliang@sciences.sdsu.edu

Received 7 August 2006

Accepted 16 September 2006

Online 14 October 2006


In the title compound, $[\text{Zn}(\text{C}_2\text{H}_3\text{N})(\text{C}_{16}\text{H}_{21}\text{N}_3\text{O})](\text{ClO}_4)_2 \cdot \text{H}_2\text{O}$, the Zn^{II} ion is coordinated by two pyridyl N atoms, one amine N atom, and an ethanol O atom from the N, N', N'', O -tetradentate 2-[bis(2-pyridylethyl)amino]ethanol donor ligand. The fifth coordination site is filled by an acetonitrile N atom, and there is one solvent water molecule in the asymmetric unit. The 2+ charge of the cationic portion of the complex is balanced by two perchlorate counter-anions.

Comment


Zinc complexes of multidentate N-donor ligands with pendant alcohol or alkoxide O-donors have been useful for modelling alkaline phosphatase, a Zn^{II} -containing phosphomonoesterase that catalyzes the hydrolysis of phosphate monoesters at high pH (Kimura *et al.*, 1994). Recently, we reported the crystal structure of a new Zn^{II} complex containing the 2-[bis(2-pyridylethyl)amino]ethanol ligand, (I), in which the ethanol group was deprotonated upon coordinating to a Zn^{II} centre to form a bridging alkoxide ligand in a dimeric Zn^{II} structure, (II), in the solid state (Zhang & Liang, 2004). The previous crystal structure was obtained by crystallizing the complex in an aqueous solution. We report here on a crystal structure of a similar complex, (III), crystallized from acetonitrile, in which the monomeric Zn^{II} complex of the ligand 2-[bis(2-pyridylethyl)amino]ethanol is obtained and the ethanol H atom remains bound to the O atom of the ethanol unit.

As shown in Fig. 1, complex (III) has five donor atoms coordinated to the Zn^{II} centre. The calculated τ value for the Zn^{II} centre in (III) is 0.23, which shows that it is closer to square-pyramidal geometry [$\text{N}2-\text{Zn}1-\text{N}4 = 159.35 (6)^\circ$ and $\text{N}1-\text{Zn}1-\text{O}1 = 145.36 (6)^\circ$; $\tau = 0$ correlates to perfect square-pyramidal geometry and $\tau = 1$ correlates to perfect trigonal-bipyramidal geometry (Addison *et al.*, 1984)]. The structure of


(III) can be viewed as having atoms N1, N2, N4, O1 and Zn1 as the basal atoms and N3 in the apical position of a pseudo-square-pyramidal complex.



The Zn–ligand bond lengths in (III) (Table 1) are similar to those found in analogous Zn^{II} complexes. For example, the Zn– N_{py} bond lengths (py is pyridine) of 2.0418 (15) Å ($\text{Zn}1-\text{N}1$) and 2.0498 (14) Å ($\text{Zn}1-\text{N}3$) are similar to the analogous Zn– N_{py} bond lengths in (II) [2.049 (4) and 2.068 (4) Å; Zhang & Liang (2004)]. However, the amine $\text{N}2-\text{Zn}1$ bond of 2.1336 (14) Å in (III) is shorter than that found in the dimeric complex [2.231 (4) Å]. The longer Zn– N_{amine} bond in (II) can be explained by the fact that the Zn^{II} centre in (II) adopts a more trigonal-pyramidal shape than in (III), with a τ value of 0.46. Because the amine atom N3 of (II) can be considered as more of an axial ligand in a trigonal-bipyramidal geometry than the analogous amine atom N2 in (III), the Zn– N_{amine} bond in (II) is expected to be longer. Comparison of the Zn–O bonds in complexes (II) and (III) also reveals differences in bond lengths [1.959 (3) Å in (II) and 2.0552 (14) Å in (III)]. This observed lengthening of the

**Figure 1**

The molecular structure of (III), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radii. Dotted lines denote possible hydrogen-bond interactions.


Figure 2

A view of the molecular packing of (III), projected perpendicular to the *a* axis, showing the hydrogen-bond network (dotted lines denote possible hydrogen-bond interactions). H atoms have been omitted for clarity. [Symmetry codes: (i) *x*, *y* − 1, *z*; (ii) 2 − *x*, 2 − *y*, 2 − *z*; (iii) 1 − *x*, 2 − *y*, 2 − *z*; (iv) 2 − *x*, 1 − *y*, 2 − *z*; (v) 1 − *x*, 1 − *y*, 2 − *z*.]

Zn—O bond in (III) is consistent with a Zn^{II} metal centre bonded to a neutral ethanol O-atom donor, whereas the intramolecular Zn—O bond in (II) involves a monoanionic alkoxide O-atom donor. Several previous studies have shown that Zn^{II}–alcohol complexes with neutral coordinating O-atom donors have Zn—O bonds longer than 2 Å (Follner, 1972; Lutz & Bakker, 2003; Iranzo *et al.*, 2003), while Zn^{II}–alkoxide compounds typically exhibit Zn—O bonds shorter than 2 Å (Kimura *et al.*, 1994; Hahn *et al.*, 2003; Davies *et al.*, 1998).

Inspection of the asymmetric unit of (III) (Fig. 1) reveals one complex cation, one uncoordinated water molecule and two perchlorate anions. The ethanol OH group of (III) forms an intermolecular O1—H1A···O10 hydrogen bond with a water molecule, which further links two adjacent perchlorate anions *via* O10—H10A(H10B)···O2(O9) close contacts (Table 2). As can be seen in Fig. 2, the organization of this motif is influenced by a complex blend of weak C—H···perchlorate interactions that extend along the *a* axis. Although a structural feature of this organization is inversion-related face-to-face stacking of pyridyl fragments, only the pyridyl (N3) moiety adopts a favourable geometry for π – π stacking [interplanar distance = 3.434 (3) Å and centroid–centroid distance = 3.862 (3) Å].

Experimental

Caution! Although no problems were encountered in this work, metal perchlorate complexes are potentially explosive and should be handled with proper precautions and in small amounts.

Complex (III) was synthesized by treating the ligand (I) (0.030 g, 0.11 mmol) with 1 equivalent of Zn(ClO₄)₂·6H₂O (0.041 g, 0.11 mmol) in CH₃CN (10 ml) with stirring for 1 h at room temperature. Diethyl ether (25 ml) was then layered on the light-yellow acetonitrile solution and the mixture was set aside for crystallization. After 4 d, colourless crystals of (III) had formed (0.037 g, 0.058 mmol, 53% yield). A single crystal from this procedure was used for X-ray analysis. ¹H NMR (500 MHz, CD₃CN): δ 8.55 (*d*, py-6H), 8.19 (*t*, py-4H), 7.71 (*d*, py-5H), 7.60 (*t*, py-3H), 3.86 (*t*, NCH₂CH₂OZn), 3.28–3.04 (py-CH₂CH₂N), 2.97 (*t*, NCH₂CH₂OZn); ¹³C NMR (CD₃CN): δ 162.1 (py-2), 149.9 (py-6), 142.1 (py-4), 127.1 (py-3), 124.7 (py-5), 60.7 (NCH₂CH₂OZn), 58.5 (NCH₂CH₂OZn), 54.5 (py-CH₂CH₂N), 33.5 (py-CH₂CH₂N); FT-IR (KBr, ν , cm^{−1}):

3448 (O—H, *str*, *br*), 3073, 2948, 2926, 2871, 2809 (C—H, *str*), 1608, 1571, 1486, 1445 (pyridine ring), 1085, 622 (ClO₄[−]). Elemental analysis calculated for C₁₈H₂₆Cl₂N₄O₁₀Zn: C 36.35, H 4.41, N 9.42%; found: C 35.98, H 4.67, N 9.05%. MS (ESI): *m/z* 334 ([L¹–H + M – CH₃CN]⁺).

Crystal data

[Zn(C₂H₃N)(C₁₆H₂₁N₃O)](ClO₄)₂·H₂O
M_r = 594.72
 Monoclinic, *P*2₁/*c*
a = 17.3792 (13) Å
b = 7.8924 (6) Å
c = 18.6583 (14) Å
 β = 109.487 (1)°
V = 2412.6 (3) Å³
Z = 4
D_x = 1.637 Mg m^{−3}
 Mo K α radiation
 μ = 1.30 mm^{−1}
T = 193 (2) K
 Block, colourless
 0.35 × 0.20 × 0.10 mm

Data collection

Bruker SMART CCD area-detector diffractometer
 φ and ω scans
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
T_{min} = 0.659, *T_{max}* = 0.881
 16837 measured reflections
 5502 independent reflections
 4695 reflections with *I* > 2 σ (*I*)
R_{int} = 0.031
 θ_{\max} = 27.5°

Refinement

Refinement on *F*²
R[*F*² > 2 σ (*F*²)] = 0.037
wR(*F*²) = 0.098
S = 1.05
 5502 reflections
 409 parameters
 H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0503P)^2 + 1.2239P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.91 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.40 \text{ e } \text{Å}^{-3}$

Table 1

Selected geometric parameters (Å, °).

Zn1—N1	2.0418 (15)	Zn1—N4	2.1273 (16)
Zn1—N3	2.0498 (14)	Zn1—N2	2.1336 (14)
Zn1—O1	2.0552 (14)		
N1—Zn1—N3	115.63 (6)	N1—Zn1—N2	95.93 (6)
N1—Zn1—O1	145.36 (6)	N3—Zn1—N2	98.17 (6)
N3—Zn1—O1	98.96 (6)	O1—Zn1—N2	80.23 (6)
N1—Zn1—N4	90.27 (6)	N4—Zn1—N2	159.35 (6)
N3—Zn1—N4	96.83 (6)	N4—C17—C18	179.7 (3)
O1—Zn1—N4	83.49 (6)		
O1—Zn1—N2—C15	22.41 (11)	N1—Zn1—N2—C7	−3.19 (12)
N3—Zn1—N2—C8	1.24 (13)		

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1A···O10	0.73 (3)	1.88 (3)	2.602 (2)	170 (3)
O10—H10A···O2	0.76 (3)	2.07 (3)	2.836 (3)	175 (3)
O10—H10B···O9	0.76 (3)	2.06 (3)	2.809 (3)	172 (3)
C1—H1···O5 ⁱ	0.94 (2)	2.57 (2)	3.318 (3)	136 (2)
C6—H6A···O3 ⁱⁱ	0.88 (3)	2.58 (2)	3.293 (3)	139 (2)
C8—H8A···O6 ⁱⁱⁱ	0.97 (2)	2.79 (2)	3.432 (3)	124 (1)
C9—H9B···O8 ⁱⁱⁱ	0.87 (2)	2.67 (2)	3.402 (3)	143 (2)
C14—H14···O7 ⁱ	0.99 (2)	2.42 (2)	3.303 (3)	149 (2)

Symmetry codes: (i) *x*, *y* − 1, *z*; (ii) −*x* + 2, −*y* + 2, −*z* + 2; (iii) −*x* + 1, −*y* + 2, −*z* + 2.

Methyl H atoms were treated as riding, with C—H = 0.98 Å and *U_{iso}*(H) = 1.5*U_{eq}*(C), and were allowed to rotate freely during refinement using the AFIX 137 command of *SHELXTL* (Bruker,

2000). All other H atoms were located in a difference density map and refined isotropically.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SMART*; data reduction: *SAINT* (Bruker, 2002) and *XPREP* (Bruker, 2001); program(s) used to solve structure: *SHELXTL* (Bruker, 2000); program(s) used to refine structure: *SHELXTL*; molecular graphics: *X-SEED*; software used to prepare material for publication: *X-SEED* (Barbour, 2001).

The authors thank San Diego State University and the San Diego Foundation Blasker–Rose–Miah Fund for financial support. We are also grateful to Dr LeRoy Lafferty for help in obtaining the NMR spectra.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FA3036). Services for accessing these data are described at the back of the journal.

References

- Addison, A. W., Rao, T. N., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1349–1356.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2000). *SADABS* (Version 2.05) and *SHELXTL* (Version 6.10). Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2001). *SMART* (Version 5.625) and *XPREP* (Version 6.12). Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2002). *SAINT*. Version 6.36a. Bruker AXS Inc., Madison, Wisconsin, USA.
- Davies, P. J., Taylor, M. R. & Wainwright, K. P. (1998). *Chem. Commun.* pp. 827–828.
- Follner, H. (1972). *Acta Cryst.* **B28**, 157–160.
- Hahn, F. E., Jocher, C., Lugger, T. & Pape, T. (2003). *Z. Anorg. Allg. Chem.* **629**, 2341–2147.
- Iranzo, O., Kovalevsky, A. Y., Morrow, J. R. & Richard, J. P. (2003). *J. Am. Chem. Soc.* **125**, 1988–1993.
- Kimura, E., Nakamura, I., Koike, T., Shionoya, M., Kodama, Y., Ikeda, T. & Shiro, M. (1994). *J. Am. Chem. Soc.* **116**, 4764–4771.
- Lutz, M. & Bakker, R. (2003). *Acta Cryst.* **E59**, m74–m76.
- Zhang, Y. & Liang, H.-C. (2004). *Inorg. Chem. Commun.* **9**, 460–463.