

January 2007

Life Without Water: Expression of Plant LEA Genes by an Anhydrobiotic Arthropod

Steven C. Hand
Louisiana State University

Dana Jones
Louisiana State University

Michael A. Menze
Eastern Illinois University, michael.menze@louisville.edu

Trudy L. Witt
Louisiana State University

Follow this and additional works at: http://thekeep.eiu.edu/bio_fac

 Part of the [Biology Commons](#)

Recommended Citation

Hand, Steven C.; Jones, Dana; Menze, Michael A.; and Witt, Trudy L., "Life Without Water: Expression of Plant LEA Genes by an Anhydrobiotic Arthropod" (2007). *Faculty Research & Creative Activity*. 29.
http://thekeep.eiu.edu/bio_fac/29

This Article is brought to you for free and open access by the Biological Sciences at The Keep. It has been accepted for inclusion in Faculty Research & Creative Activity by an authorized administrator of The Keep. For more information, please contact tabruns@eiu.edu.

Life Without Water: Expression of Plant LEA Genes by an Anhydrobiotic Arthropod

STEVEN C. HAND*, DANA JONES, MICHAEL A. MENZE,
AND TRUDY L. WITT

*Division of Cellular, Developmental, and Integrative Biology, Department of
Biological Sciences, Louisiana State University, Baton Rouge, Louisiana 70803*

ABSTRACT Anhydrobiotic animals protect cellular architecture and metabolic machinery in the dry state, yet the molecular repertoire supporting this profound dehydration tolerance is not fully understood. For the desiccation-tolerant crustacean, *Artemia franciscana*, we report differential expression of two distinct mRNAs encoding for proteins that share sequence similarities and structural features with late-embryogenesis abundant (LEA) proteins originally discovered in plants. Bioinformatic analyses support assignment of the LEA proteins from *A. franciscana* to group 3. This eucoelomate species is the most highly evolved animal for which LEA gene expression has been reported. It is becoming clear that an ensemble of micromolecules and macromolecules is important for establishing the physical conditions required for cellular stabilization during drying in nature. *J. Exp. Zool.* 307A:62–66, 2007. © 2006 Wiley-Liss, Inc.

How to cite this article: Hand SC, Jones D, Menze MA, Witt TL. 2007. Life without water: expression of plant lea genes by an anhydrobiotic arthropod. *J. Exp. Zool.* 307A: 62–66.

It is a biological truism that environmental stresses that impact the activity of cellular water pose a threat to life (Somero, '92). For centuries scientists have been intrigued by exceptional animals that can survive the loss of virtually all cellular water for prolonged periods (Leeuwenhoek, 1702; Crowe and Clegg, '73). The mechanisms by which these anhydrobiotic animals protect cellular architecture and metabolic machinery in the dry state are not only of biological interest but also of biomedical importance for cell stabilization (Crowe et al., 2005; Elliott et al., 2006). For a desiccation-tolerant arthropod, we report herein the differential expression of two distinct mRNAs encoding for proteins that share strong sequence similarities and structural features with late-embryogenesis abundant (LEA) proteins originally discovered in plants.

Intracellular sugar glasses like those formed with trehalose or other carbohydrates during drying (Hoekstra, 2005 and references therein) provide protection during water stress in animals and plants. Hydrophilic LEA proteins (hydrophilins) seen for example in seeds, pollen, and the resurrection plant are receiving considerable interest because their intracellular accumulation

is tightly correlated with acquisition of desiccation tolerance, and recent in vitro data indicate their capacity to stabilize other proteins during drying (Grelet et al., 2005). Embryos of the brine shrimp *Artemia franciscana* have served as an important model species for studies of animal desiccation, and evidence from this organism has underscored the role of trehalose and small stress proteins in anhydrobiosis and other environmental stresses (Crowe et al., '84; Warner et al., 2004; Clegg, 2005; Crowe et al., 2005). In this study, we investigated whether or not expression of other desiccation-associated proteins might be found in the developmental stages of the brine shrimp that exhibit survival in the dried state.

Grant sponsor: NIH Grant; Grant number: 1-RO1-GM0-71345-01 (General Medical Sciences).

*Correspondence to: S.C. Hand, Department of Biological Sciences, 202 Life Sciences Building, Louisiana State University, Baton Rouge, LA 70803. E-mail: shand@LSU.edu

Received 20 June 2006; Accepted 8 August 2006

Published online 15 November 2006 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/jez.a.343.

MATERIAL AND METHODS

Animals

Encysted embryos of the brine shrimp, *A. franciscana* Kellog (Great Salt Lake population), were either obtained in the dehydrated state (post-diapause) from Sanders Brine Shrimp Co. (Ogden, UT) or collected from the surface of the Great Salt Lake in the hydrated state (diapause) (Covi and Hand, 2005). Embryo viability and preservation of the diapause state during storage were evaluated as previously described (Reynolds and Hand, 2004).

Sequencing of cloned LEA genes

DNA inserts were sequenced directly from plasmids isolated from bacterial clones picked from a full-length unidirectional cDNA Library (Lambda Uni-ZAP XR Vector). The library was prepared from poly-A mRNA purified and pooled from active and diapause stage embryos. Sequencing utilized BigDye terminator chemistry and an ABI PRISM 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA). Primer walking was used to insure full-length sequence was obtained. Sequences were assembled using Sequencher software (Gene Codes Co., Ann Arbor, MI).

Isolation of total RNA and preparation of cDNA for quantitative PCR

Post-diapause embryos were hydrated and incubated at room temperature in medium equilibrated with air as described previously (Covi and Hand, 2005). Briefly, after the desired time periods of incubation, RNA extractions were performed with an RNeasy Midi kit (Qiagen, Valencia, CA) as per the manufacturer's instructions for animal tissues. For samples of larvae (24 hr), swimming nauplii were separated from the shed chorion. Diapause embryos were incubated for 4 d as previously reported (Reynolds and Hand, 2004) to allow individuals not in diapause to hatch; larvae and empty shells were removed. The concentration of RNA in each sample was determined spectrophotometrically (A_{260}). AMV reverse transcriptase (Promega, Madison, WI) and a dT₂₅V primer were employed to reverse transcribe poly(A⁺) mRNA from total RNA.

Real-time quantitative PCR

Analyses of cDNA from various developmental stages of *A. franciscana* were performed using

real-time quantitative polymerase chain reaction (rtqPCR) with an iCycler (Bio-Rad Laboratories, Hercules, CA) and SYBR Green. PCR reaction mixes were prepared by using the IQ SYBR Green Supermix (Bio-Rad Laboratories) with a total reaction volume of 25 μ l. Cycling parameters were 3 min at 95°C and then 50 cycles at 95°C (15 sec) and 56°C (30 sec), followed by a melting curve analysis. Each developmental stage was evaluated with 4–6 independent experiments, each with three nested replicates, and with reference dye normalization. The cycle threshold value (Ct) determined with the manufacturer's software and used for analyses, and all Δ Ct values were normalized to the expression of a bona fide housekeeping gene, α -tubulin (Zheng et al., '98), after determination of the PCR efficiencies (Pfaffl, 2001). Primer sequences for α -tubulin were 5'-CGA CCA TAA AAG CGC AGT CA-3' and 5'-CTA CCC AGC ACC ACA GGT CTC T-3'. Primer sequences for *Afrlea1* were 5'-GTG CCG TCT GTG CTC TC-3' and 5'-CAG GGA GCC TAT GAG GGA CT-3'. Primer sequences for *Afrlea2* were 5'-GAA TGT GCA GCA TCA GCA GT-3' and 5'-GCT CAG TCA ACA TAT GAC CCA GTG-3'. Values for the fold change in expression are given as means \pm SD, and significance was assessed by applying the *t*-test of the mean with Minitab 1998 (Minitab Inc., State Collage, PA).

Bioinformatic analyses

Sequences were compared with the GenBank/NCBI database using BLAST software, and also to the Pfam database (www.sanger.ac.uk/Software/Pfam/search.shtml). The Kyte and Doolittle algorithm (Kyte and Doolittle, '82) was used to construct hydropathy plots (ProtScale program, <http://au.expasy.org/tools/protscale.html>). A 9-residue moving average was used to compute the hydropathy score. The Biology Workbench 3.2 site (<http://workbench.sdsc.edu/>) was used for multiple sequence alignments (CLUSTALW) and evaluation of amino acid abundance (AASTATS). Secondary structure predictions were run with the GOR IV, PHDsec, HNN, SSpro, SOPMA, Porter, and Prof programs (<http://au.expasy.org/tools/>), and values for alpha-helix content were averaged. Subcellular localization of proteins was predicted using: PREDOTAR (www.inra.fr/predotar/), TargetP (www.cbs.dtu.dk/services/TargetP/), and MitoProt (<http://ihg.gsf.de/ihg/mitoprot.html>). Both proteins were evaluated for regions that could be involved in coiled-coil structures using the

program COILS (www.ch.embnet.org/software/COILS_form.html). Multiple sites for coiled-coil structures were identified for AfrLEA1 (at least 6) and AfrLEA2 (at least 3).

RESULTS AND ANALYSIS

Based on data mining of a cDNA library prepared from *A. franciscana* embryos, coupled with real-time quantitative PCR (rtqPCR), we find that each developmental stage with the capacity for anhydrobiosis (diapause and post-diapause embryos) expresses high levels of two LEA mRNAs, while the desiccation-intolerant larval stage (control) shows expression that is many fold lower (Fig. 1A). This differential expression is consistent with a role for these gene products in survivorship during dehydration.

Deduced protein sequences indicate that AfrLEA1 is composed of 357 amino acids with an apparent molecular mass of 39 kDa, while AfrLEA2 contains 364 amino acids and is 39 kDa.

(Table 1). Both proteins are strongly hydrophilic based on hydropathy plots (Fig. 1B). Further, the hydropathy patterns clearly reveal repeating motifs characteristic of plant LEA proteins. Sequence analyses show conserved repeats of 32 amino acids for AfrLEA1 and 14 for AfrLEA2; the tandem repeats become more degenerate toward the carboxy terminus (Fig. 1C, D). The repeating motifs do not match either the historical 11-mer repeat seen for some group 3 LEA proteins (Dure, '93; Cuming, '99), or the 20-mer sometimes seen for group 1 LEA proteins (Esperlund et al., '92; Cuming, '99). However, lack of concordance with the historical repeats is observed even for plant LEA proteins (cf. Grelet et al., 2005).

Comparisons with the GenBank database reveal strong similarities to a phylogenetically broad suite of group 3 LEA proteins for AfrLEA1 (e.g., accession numbers: NP_001024042, *Caenorhabditis elegans*, nematode, E value = $5e^{-15}$; AAA85367, *Picea glauca*, white spruce, $1e^{-08}$; CAA 80491, *Glycine max*, soybean, $1e^{-06}$; YP604937,




Fig. 1. (A) mRNA expression profiles for the LEA genes *Afrlea1* and *Afrlea2* from *Artemia franciscana* embryos. LEA mRNAs are maintained 7–14-fold higher in the two desiccation-tolerant embryonic stages (i.e., diapause and post-diapause) compared to the desiccation-intolerant nauplius larva that served as a control (mean \pm SD; $N = 4-5$; $P < 0.005$). The double asterisks indicate that the paired means are statistically different. All Δ Ct values (corrected for efficiency) were normalized to the expression of a housekeeping gene (α -tubulin). (B) Kyte and Doolittle hydropathy plots for the deduced proteins AfrLEA1 and AfrLEA2 indicate strong hydrophilicity, as shown by values below zero. Amino acid position is plotted on the x-axis beginning with the N-terminus. The more evident cyclic repeats are numbered sequentially above the x-axis for each protein (color coded). Alignments of tandemly repeated motifs, which correspond to the numbered hydropathy patterns, are provided in (C) and (D). The repeats become more degenerate as one moves toward the C-terminus.

TABLE 1. Deduced sequences for LEA proteins AfrLEA1 and AfrLEA2 from *Artemia franciscana* embryos

AfrLEA1						
MAEPEEPPGI	YEKVKSAFVS	APDRAQEAYN	QAYESARSVF	DDAVRSARKM	KNTAAEQAQQ	60
AYEGLKESPE	NLQRVTRDIY	HQAQDTGKGA	YETVAGSADD	AYRRAQETAQ	AAQEQSKGFL	120
NRVKDTLTAP	FSSSSDQAKE	TYDRTKDEAQ	YRAQQAADAG	QGFFGKVKDT	ITAPFTSGYD	180
QTQEGYERAR	RSAEAAAQQA	ADQGGTLFER	AKDTITSPFS	SGSEQAQESF	ERAKRAAEEQ	240
VEQSKGMFQN	IKGTITSPFN	SAADTAKEAG	QRAKKQAEAA	ADQSQGFMMQK	VKDTVASPFLL	300
SAGEESQEI	ERTKREAEAA	RHQGEGFLHR	VADTIMHPFQ	SSSEQVGEAA	DRIKRGAA	357
AfrLEA2						
MPKAAAKGIG	ETVKADADV	EGMASTGYEK	LKSAFGIASN	KTKDAAENVA	ESARATKDYT	60
VDSAKSAYDK	TVDSTKSAYD	KTTDSAKSVH	DSTADTAKSA	YNKATETLGS	AYDKTKDTAQ	120
STYDQVTGAA	HSAYDKTAEA	TKSAYDKTAD	AAHSVYNKGT	DAGKQAYDST	KEAARSTGKS	180
ISDAAYFTGK	GAERQGDQVK	SELPSYSPSS	SKEKLAQHLV	KSEKEGKCLT	EEALKDRDLS	240
QVPGFRSVK	AHEPDAKEDI	SAVDFASASP	SQRKVADTEG	VWSSPVDRQE	SRFFSDLAGK	300
IGDMLGGGKI	NAIQTPPEMD	HERLIHKSSQ	SQVAGNVPR	AKTAWTPEDR	IILHQRFPK	360
ENPE						364

Deinococcus radiodurans, non-spore-forming bacterium, $9e^{-05}$) and for AfrLEA2 (e.g., BAA11017, *Arabidopsis thaliana*, thale cress, $2e^{-14}$; AD59387, *Brassica napus*, rapeseed, $5e^{-15}$; ABA26579, *Phaseolus vulgaris*, bean, $7e^{-13}$; CAF32327, *Pisum sativum*, pea, $3e^{-11}$). Wise (2003) has emphasized that in addition to strong sequence similarities to canonical LEA proteins, upregulated expression profiles that are tightly associated with desiccation, as we report here, are also important for LEA protein identification. In this context, a LEA-like DNA sequence from a chironomid insect larva is present in GeneBank (accession number BAE92616), but expression data are unavailable.

Other features consistent with assignment of AfrLEA1 and AfrLEA2 to group 3 are: high alpha-helix content (77%, AfrLEA1; 59%, AfrLEA2) with significant potential for coiled-coil structures, over-representation of alanine (16.2%, AfrLEA1; 14.3%, AfrLEA2), and the lack of over-representation for glycine or arginine. Both sequences are devoid of cysteine, a feature of many LEA proteins. Finally, when the two *A. franciscana* sequences were compared against the Pfam database, both were assigned to Pfam LEA 4 family, which corresponds to group 3 LEA proteins (cf. Grelet et al., 2005).

DISCUSSION

To our knowledge, this is the first report of two LEA proteins expressed in one animal species. Targeting to different cellular locations might explain the functional significance of two LEA proteins, although bioinformatic analyses with

subcellular targeting programs failed to suggest localization to specific organelles such as the mitochondrion. The predictive power of these current software programs is limited, and the complexity of the N-terminal cleavable extensions (termed presequences) involved in targeting is high (Pfanner and Geissler, 2001; Wiedemann et al., 2004).

The expression of LEA proteins is not restricted to plants, having been documented in bacteria, fungi, nematodes (Wise and Tunnacliffe, 2004; Hoekstra, 2005, and references therein), and now a desiccation-tolerant crustacean, *A. franciscana*. This eucoelomate species is the most highly evolved animal for which LEA gene expression has been reported. Among various physiological roles, stabilization of sugar glasses is often suggested for LEA proteins (Hoekstra, 2005), along with protein stabilization via protein-protein interaction (Grelet et al., 2005), ion sequestration (Grelet et al., 2005), and formation of structural networks (Wise and Tunnacliffe, 2004). Such networks have been hypothesized to increase cellular resistance to physical stresses imposed by desiccation (cf. Goyal et al., 2003). It is becoming clear that an ensemble of micromolecules and macromolecules is important for establishing the physical conditions required for cellular stabilization during drying in nature.

LITERATURE CITED

- Clegg JS. 2005. Desiccation tolerance in encysted embryos of the animal extremophile, *Artemia*. *Integr Comp Biol* 45: 715–724.
- Covi JA, Hand SC. 2005. V-ATPase expression during development of *Artemia franciscana* embryos: potential role

- proton gradients in anoxia signaling. *J Exp Biol* 208: 2783–2798.
- Crowe JH, Clegg JS, editors. 1973. *Anhydrobiosis*. Stroudsburg, PA: Dowden Hutchinson Ross. 477p.
- Crowe JH, Crowe LM, Chapman D. 1984. Preservation of membranes in anhydrobiotic organisms—the role of trehalose. *Science* 223:701–703.
- Crowe JH, Crowe LM, Wolkers WF, Oliver AE, Ma X, Auh J-H, Tang M, Zhu S, Norris J, Tablin F. 2005. Stabilization of dry mammalian cells: lessons from nature. *Integr Comp Biol* 45:810–820.
- Cuming AC. 1999. LEA proteins. In: Shewry P, Casey R, editors. *Seed proteins*. Boston: Kluwer Academic Publishers. p 753–780.
- Dure L III. 1993. A repeating 11-mer amino acid motif and plant desiccation. *Plant J* 3:363–369.
- Elliott G, Liu X-H, Cusick JL, Menze MA, Vincent J, Witt T, Hand SC, Toner M. 2006. Trehalose uptake through P2X₇ purinergic channels provides dehydration protection. *Cryobiology* 52:114–127.
- Esperlund M, Saeboe-Larssen S, Hughes DW, Galau GA, Larsen F, Jakobsen KS. 1992. Late embryogenesis-abundant genes encoding proteins with different numbers of hydrophilic repeats are regulated differentially by abscisic acid and osmotic stress. *Plant J* 2:241–252.
- Goyal K, Tisi L, Basran A, Browne J, Burnell A, Zurdo J, Tunnacliffe A. 2003. Transition from natively unfolded to folded state induced by desiccation in an anhydrobiotic nematode protein. *J Biol Chem* 278:12977–12984.
- Grelet J, Benamar A, Teyssier E, Avelange-Macherel M-H, Grunwald D, Macherel D. 2005. Identification in pea seed mitochondria of a late-embryogenesis abundant protein able to protect enzymes from drying. *Plant Physiol* 137:157–167.
- Hoekstra FA. 2005. Differential longevities in desiccated anhydrobiotic plant systems. *Integr Comp Biol* 45:725–733.
- Kyte J, Doolittle RF. 1982. A simple method for displaying the hydropathic character of a protein. *J Mol Biol* 157:105–132.
- van Leeuwenhoek A. 1702. On certain animalcules found in the sediment in gutters of the roofs of houses. In: Hoole S, translator. *The select works of Antony van Leeuwenhoek*. London, 1798, 1807, Vol. 2, Letter 144. p 207–213.
- Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucl Acid Res* 29: 2002–2007.
- Pfanner N, Geissler A. 2001. Versatility of the mitochondrial protein import machinery. *Nat Rev Mol Cell Biol* 2: 339–349.
- Reynolds JA, Hand SC. 2004. Differences in isolated mitochondria are insufficient to account for respiratory depression during diapause in *Artemia franciscana* embryos. *Physiol Biochem Zool* 77:366–377.
- Somero GN. 1992. Adapting to water stress: convergence on common solutes. In: Somero GN, Osmond CB, Bolis CL, editors. *Water and life*. Berlin: Springer-Verlag. p 3–18.
- Warner AH, Brunet RT, MacRae TH, Clegg JS. 2004. Artemin is an RNA-binding protein with high thermal stability and potential RNA chaperone activity. *Arch Biochem Biophys* 424:189–200.
- Wiedemann N, Frazier AE, Pfanner N. 2004. The protein import machinery of mitochondria. *J Biol Chem* 279: 14473–14476.
- Wise MJ. 2003. LEAping to conclusions: a computational reanalysis of late embryogenesis abundant proteins and their possible roles. *BMC Bioinform* 4:52–71.
- Wise MJ, Tunnacliffe A. 2004. POPP the question: what do LEA proteins do? *Trends Plant Sci* 9:13–17.
- Zheng Y, Roy PJ, Liang P, McRae TH. 1998. Cloning and sequencing of an alpha-tubulin cDNA from *Artemia franciscana*: evidence for translational regulation of alpha-tubulin synthesis. *Biochim Biophys Acta* 1442:419–426.