

1973

Parasites from Gizzard Shad, *Dorosoma cepedianum*, in Lake Charleston, Illinois

Francis M. Yabai Kaikumba

Eastern Illinois University

This research is a product of the graduate program in [Zoology](#) at Eastern Illinois University. [Find out more](#) about the program.

Recommended Citation

Kaikumba, Francis M. Yabai, "Parasites from Gizzard Shad, *Dorosoma cepedianum*, in Lake Charleston, Illinois" (1973). *Masters Theses*. 3835.

<https://thekeep.eiu.edu/theses/3835>

This is brought to you for free and open access by the Student Theses & Publications at The Keep. It has been accepted for inclusion in Masters Theses by an authorized administrator of The Keep. For more information, please contact tabruns@eiu.edu.

PAPER CERTIFICATE #2

TO: Graduate Degree Candidates who have written formal theses.

SUBJECT: Permission to reproduce theses.

The University Library is receiving a number of requests from other institutions asking permission to reproduce dissertations for inclusion in their library holdings. Although no copyright laws are involved, we feel that professional courtesy demands that permission be obtained from the author before we allow theses to be copied.

Please sign one of the following statements:

Booth Library of Eastern Illinois University has my permission to lend my thesis to a reputable college or university for the purpose of copying it for inclusion in that institution's library or research holdings.

10 SEPTEMBER 1973

Date

I respectfully request Booth Library of Eastern Illinois University not allow my thesis be reproduced because _____

_____ Date

_____ Author

pdm

PARASITES FROM GIZZARD SHAD, DOROSOMA CEPEDIANUM,

IN LAKE CHARLESTON, ILLINOIS

(TITLE)

BY

FRANCIS M. YABAI KAIKUMBA

B.A. BEREA COLLEGE

THESIS

SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF

MASTER OF SCIENCE

IN THE GRADUATE SCHOOL, EASTERN ILLINOIS UNIVERSITY
CHARLESTON, ILLINOIS

1973

YEAR

I HEREBY RECOMMEND THIS THESIS BE ACCEPTED AS FULFILLING
THIS PART OF THE GRADUATE DEGREE CITED ABOVE

6 SEPT. 1973
DATE

6 Sept. 1973
DATE

The undersigned, appointed by the Head of the Department of Zoology,
have examined a thesis entitled

PARASITES FROM GIZZARD SHAD, DOROSOMA CEPEDIANUM,
IN LAKE CHARLESTON, ILLINOIS

Presented by

FRANCIS M. YABAI KAIKUMBA

a candidate for the degree of Master of Science
and hereby certify that in their opinion it is acceptable.

6 SEPTEMBER 1972

ACKNOWLEDGMENT

The author owes an invaluable gratitude to Dr. Bill T. Ridgeway whose incessant help and advice contributed largely to the success of this work. Fish examined were obtained through the kind help of Dr. L. Durham. Last but not least, to my loving wife Nemata, I dedicate this thesis with sincere appreciation for her encouragement and support throughout my graduate study.

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGMENT	i
LIST OF TABLES	iii
ABSTRACT	1
INTRODUCTION	1
METHODS	2
RESULTS	3
DISCUSSION	6
BIBLIOGRAPHY	13

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Parasites recovered from 80 gizzard shad (<u>Dorosoma cepedianum</u>) in Lake Charleston, Charleston, Illinois, 1972	5
2	Parasites recovered from gizzard shad (<u>Dorosoma cepedianum</u>) in U.S.A. from 1913 to 1967	11

Parasites from Gizzard Shad, Dorosoma cepedianum,
in Lake Charleston, Illinois

Abstract: Eighty gizzard shad (Dorosoma cepedianum) from Lake Charleston, Charleston, Illinois were examined for parasites. Eight fish were parasitized. Three fish contained the acanthocephalan Tanaorhampus longirostris, three fish harbored cestodes of the genus Glaridacris and two fish had the nematode Camallanus oxycephalus. The latter is here reported for the first time from gizzard shad. Twenty fish were infected with the fungus Saprolegnia sp. The numbers of helminth parasites harbored by shad are less than reported for other forage fish. However, the incident of parasitism reported in this study is greater than that reported by previous workers.

It is important to understand the ecological relationships of valued game fishes and the fish serving as food for such game fishes. Studies of the parasites in a fish species may contribute to an understanding of its ecology because parasites may be closely related to a host's feeding habits and habitat.

In lakes and streams, the gizzard shad, Dorosoma cepedianum, is utilized as food for largemouth bass, black crappie, and walleye (Kutkuhn, 1958). No study of the parasites of this major forage fish in Illinois has been reported. Therefore, this study of the parasites of the gizzard shad in Lake Charleston, Coles County, Illinois was

undertaken. Lake Charleston is a four hundred-acre lake on the Embarrass River, serving as a city water supply and for recreation.

METHODS

Fish were collected by electrofishing. Each fish was weighed and classified adult or juvenile according to criteria established by Miller (1960). Sex was determined after examining the reproductive organs. Immediately following collection pithed fish were examined for ectoparasites using a dissection microscope to scan the body surface, gills, and scrapings from the body, mouth and anus.

Oral tissue scrapings and mucous scrapings from areas of apparent injury were put into jars containing about 50 to 70 mls. of physiological saline and well shaken. The mixture was allowed to decant for ten minutes and the sediment examined for parasites. Gills of all the fish were removed and placed in physiological saline and treated in the same manner as for mucous scrapings.

The body cavities of the hosts were opened and examined for cysts and helminths. The gastro-intestinal tract was removed, cut in three sections, opened, and scanned for endoparasites. Gut contents and mucosal scrapings were added to a jar of physiological saline and the mixture treated as for scrapings and gills.

All helminths, except acanthocephalans, were killed and preserved in 70% ethanol with 2% glycerine. Acanthocephalans were placed in distilled water until the hook-bearing proboscis was extruded, killed in warm (70°C) formalin and stored in 70% ethanol.

Nematodes and acanthocephalans were examined in lactophenol on microslides. Standard staining techniques were used for cestodes.

Helminths were identified using Yamaguti (1959, 1961, 1963), Hoffman (1967), and Yorke and Maplestone (1962). Fungal mycelia in mucous scrapings were identified according to keys in Hoffman (1967).

RESULTS

Eighty gizzard shad were collected and examined for parasites. One species of nematode, one species of acanthocephalid, and one species of cestode worm were found. One kind of fungus was observed. No trematode or protozoan parasites were recovered.

The nematode, Camallanus oxycephalus, Ward and Magath, 1916, was present in the small intestine of two female fish. In both cases, only one worm (female) per host was recovered. The nematode is red with a slitlike mouth that lacks lips. A buccal capsule is present with two lateral shell-shaped chitinous valves. The valves have on their inner surface, parallel longitudinal ribs. A chitinous trident shaped supporting process is located at the point where the valves meet. The buccal capsule is surrounded by a chitinous circular collar at its entry into the esophagus. The latter has a short muscular anterior region and a long posterior region that is enlarged at its end. The length of Camallanus oxycephalus is 10 mm.

Acanthocephalan worms identified as Tanaorhamphus longirostris (Van Cleave, 1921) occurred in the upper one-third of the small intestine of three hosts. These parasites were firmly imbedded in the intestinal mucosa by their spiny proboscises. The number of parasites per infected host was one and all were females. Tanaorhamphus longirostris is a medium-sized acanthocephalan, and has a sub-cylindrical trunk with the hypodermic giant nuclei mostly in the mid dorsal line.

It has a fairly long and cylindrical proboscis armed with sixteen rows of ten hooks each. The proboscis has a ganglion at its base and the long slender lemnisci have very few nuclei. The specimens are female and the eggs are elongate oval.

Specimens of the cestode worm obtained in this study were identified only as Glaridacris sp. Cooper, 1920. All worms are monozoic and about 2 mm long. The well-defined scolex has six bothria-like depressions which are not very evident. The uterus does not extend anterior to the cirrus pouch, and the latter opens ventrally into the atrium. The H-shaped ovary is medullary. The worms occurred in the lower end of the intestine of the fish.

On twenty fish, cottony masses of fungal mycelia were observed on areas of apparent abrasion. A detailed identification of this fungus down to species level was not made. But under low magnification the fungus was observed to consist of a mass of non-septate filaments each of which was 20 μ in diameter. The filaments of older infections terminated in club-like sporangial enlargements. The fungus is Saprolegnia.

Results are tabulated in Table 1.

Table 1. Parasites recovered from 80 gizzard shad (Dorosoma cepedianum) in Lake Charleston, Charleston, Illinois, 1972.

	% of infected hosts	No. of infected hosts	No. of parasites found	Location	Age of host
NEMATODA					
<u>Camallanus oxycephalus</u>	2.5	2	2	Intestine	Adult
ACANTHOCEPHALA					
<u>Tanaorhamphus longirostris</u>	3.8	3	3	Intestine	Adult
CESTODA					
<u>Glaridacris</u> sp.	3.8	3	3	Intestine	Adult
PHYCOMYCETES					
<u>Saprolegnia</u> sp.	25.0	20	-	Body surface	Adult

DISCUSSION

Compared to other species of fish that have been studied, the gizzard shad has a low incidence of parasitism. Ward (1912) examined four individuals and found one cestode, one acanthocephala and no nematodes. Bangham and Hunter (1939) examined five specimens and found only one unidentified larval nematode. Van Cleave (1919), in a search for acanthocephala from the Illinois River, recovered only two species of helminths, Tanaorhamphus longirostris, and Gracilisentis gracilisentis from 164 gizzard shad.

Essex and Hunter (1926) found no helminths in 107 gizzard shad from the Rock and Mississippi rivers. According to the authors, "The feeding habits of the shad are such as to preclude the acquiring of an extensive parasitic fauna. Since it feeds largely on vegetable debris at the bottom of streams and lakes, very seldom ingesting animal food, there is little opportunity for parasites which have an extremely complex life cycle, to find it a suitable host."

Miller (1960) remarked that "the species (Dorosoma cepedianum) appears to be unusually free from parasites from which it is undoubtedly protected by its herbivorous food habits." Results of the work done by Jester and Jensen (1972) support the contentions of Miller (1960), and Essex and Hunter (1926). Jester and Jensen studied the life history and ecology of gizzard shad in the Elephant Butte Lake in New Mexico and found one ectoparasite, Argulus sp., on three fish and an acanthocephalan, Gracilisentis sp., from one fish. They noted that shad from the lake were host to water-fungus, Saprolegnia parasitica.

The results of this study support the contentions of authors cited above. The total incidence of infection by helminths is 10% and fungal

infection is 25%. Such figures are low compared to those obtained from most other fishes (Essex and Hunter, 1926). However, Miller's view that such a low parasitic infection is due to the herbivorous food habits of the fish cannot be supported here because gizzard shad feed on both zooplankton and phytoplankton. The preponderance of either of the two depends on availability (Jester and Jensen, 1972).

Bangham (1941b) stated that the young shad in Buckeye Lake, Ohio, often bear a heavy infection of myxosporidians. Ten out of twelve young fish collected at the lake by Bangham were infected whereas all adult individuals were free of myxosporidian cysts. Careful examination of the fish collected in the present study revealed no such cysts. It should be observed that all the fish were adults.

Myxosporidian cysts are usually near the surface of the host's skin. With the lapse of time and as the host develops, the cysts break thus releasing the spores. If not, the spores remain in the body until another fish feeds on the infected host. Young shad are often fed upon by game fish. Thus the infected young shad may be fed upon before adulthood. Those that survive may get rid of their infection as the cysts break, before adulthood. Table 2 summarizes all the helminth parasites recovered from gizzard shad from 1913 to 1967.

Camallanus oxycephalus reported in this study has been reported from several fishes in North America -- black and white crappies, bluegill, largemouth bass, yellow perch, sauger, white bass, northern pike, channel catfish, paddle fish, bowfin and brown trout (Hoffman, 1967). But it has never been reported from the gizzard shad.

Baer (1951), outlined the life history of Camallanus as including an egg stage, three successive larval stages (L₁, L₂, L₃), and an adult

stage. The first stage larva develops in the uterus and is consequently eliminated into the water. It is swallowed by a copepod and passed down into the body cavity of the latter. Here it further develops into the L₂ stage through the L₃ stage. If an infected copepod is eaten by a fish or any other predator the larva escapes into the latter and becomes an adult.

According to Dogiel (1958), the life history of Camallanus utilizes a crustacean intermediate host and a carrier host. The latter corresponds to what Baer (1951) calls the paratenic host. The latter host may be optional. McDaniel (1963) found that the carrier host of the parasite from turtles could be damselfly nymphs, dragonfly nymphs, or even young fish. It is possible that Camallanus oxycephalus utilizes a similar carrier or paratenic host.

Since the gizzard shad is a plankton feeder (Jester and Jensen, 1972), specimens of the fish found to be infected with C. oxycephalus, in this study, may have obtained the parasite through feeding on infected copepods. Infection can occur in both young and adult shad. Accepting Baer's statement (1951) that the carrier host is optional, it is possible that such a host is by-passed in this instance. Dogiel (1958) also believed that the viviparous Camallanus can develop without a carrier host.

There is a possibility also that the gizzard shad may have obtained the parasite through a carrier host, like a damselfly nymph, or a dragonfly nymph, or even a young fish as McDaniel (1963) stated. The shad is known to migrate periodically from the top to the bottom zones of lakes and streams (Jester and Jensen, 1972). As it dredges along the bottom of the lake, it leaves its mouth open. In this way several unavoidable materials including damselfly and dragonfly nymphs are passively ingested.

This is evidenced by the fact that the author often encountered skeletal remains of such insect nymphs in the gut contents of the fish.

Tanaorhamphus longirostris was reported from Dorosoma cepedianum in the Illinois River by Van Cleave in 1919. The acanthocephalan, according to Van Cleave, showed a strong evidence of restriction to the gizzard shad from that region. Bangham (1942) found the same parasite in the Dorosoma cepedianum, Ictalurus anguila, and Ictalurus punctatus in Tennessee.

Compared to the 75.4% infection reported in Van Cleave's work (1919), results in the present study show only 3.8%. However since gizzard shad secures its acanthocephala only through feeding upon infected copepods, the degree to which the fish is parasitised depends on the number of infected copepods eaten. Also, according to Van Cleave, seasonal conditions influence infection.

Glaridacris is a caryophyllid tapeworm that is common among suckers (Hoffman, 1967). A species, G. confusus was recovered from Dorosoma cepedianum by Fischthal (1947, 1950) in Wisconsin. It is possible that the three specimens found in the present study are G. confusus. The parasite inhabits the intestine of vertebrate hosts and tubericid oligochaetes serve as intermediate hosts.

Dogiel (1958) stated that the abundance of a host influences number and diversity of parasites. Due to its high reproductive potential and rapid growth rate, the gizzard shad is known for its tendency to over-populate waters. In this regard, the fish is expected to harbor a great abundance and variety of parasites because the latter may have the opportunity to spread from one host to another. Results in this study and others negate this view. Despite high population numbers, gizzard shad from Lake Charleston show the general freedom from parasites reported by other workers.

In conclusion, to clarify this situation, it is recommended that some careful investigation be made into the receptiveness of gizzard shad to various parasites. Such a work may somehow answer the question of low parasitic incidence and also throw light on the ecological status of the fish.

Table 2. Parasites recovered from gizzard shad (Dorosoma cepedianum) in U.S.A. from 1913 to 1967.

Species	Location in host	Type of report and geographical distribution	Author
PROTOZOA			
<u>Plistophora cepedianae</u>	Viscera	Case report Ohio	Putz, Hoffman & Dunbar, 1965
TREMATODA			
<u>Clinostomum</u> sp.	Muscles & skin	General report U.S.A. & Asia	Leidy, 1856 Bangham, 1951, 1955 Fischthal, 1945, 1950
<u>Diplostomulum</u> sp.	Muscles	Case report No. American & Europe	Hoffman, 1960
<u>Mazocracoides megalocotyle</u>	Gills	Case report Tennessee, U.S.A.	Price, 1958
<u>Mazocracoides oletangienses</u>	Gills	Case report Ohio, Tennessee, U.S.A.	Sroufe, 1958
<u>Mazocracoides similis</u>	Gills	Case report Ohio, Tennessee, U.S.A.	Price, 1959
CESTODA			
<u>Glaridacris confusus</u>	Intestine	Case report Mississippi, Iowa	Hunter, 1927
<u>Proteocephalus</u> sp.	Intestine	General report U.S.A.	Hoffman, 1967

Table 2. (Continued)

Species	Location in host	Type of report and geographical distribution	Author
ACANTHOCEPHALA			
<u>Gracilisentis gracilisentis</u>	Intestine	Case report Illinois, Mississippi	Van Cleave, 1919, 1931
<u>Tanaorhamphus longirostris</u>	Intestine	Case report Illinois, Tennessee	Van Cleave, 1919 Bangham, 1942
<u>Neoechinorhynchus longirostris</u>	Intestine	Case report Mississippi	Van Cleave, 1931
CRUSTACEA			
<u>Argulus alosae</u>	Gills, Body surface	General report U.S.A.	Hoffman, 1967
<u>Argulus appendiculosus</u>	Gills, Body surface	General report U.S.A.	Hoffman, 1967
<u>Ergasilus lanciolatus</u>	Gills, Body surface	Case report Kentucky	Wilson, 1916

BIBLIOGRAPHY

- Adams, J. R. 1958. A parasitic copepod (Selmincola falculata) attached to a fish heart. *J. Parasit.* 42(3):296.
- Alexander, C. G. 1951. A new species of Proteocephalus (cestoda) from Oregon trout. *J. Parasit.* 37(2):150-184.
- _____. 1960. A survey of parasites of Oregon trout. Rep. to Oregon State Game Comm. 1-34.
- Ali, S. M. 1956. Studies on the nematode parasites of fishes and birds found in Hyderabad state. *Indian J. Helminth.* 8(1):1-83.
- _____. 1960. On two new species of Procamallanus Baylis, 1923 from India, with a key to the species. *J. Helminth.* 34(162):12-138.
- Allison, L. N. 1950. Common diseases of fish in Michigan. Michigan Dept. Conserv., Misc. Publ. no. 5:1-27.
- Arnold, J. G., Jr. 1933. Some trematodes of the common bullhead (Ameiurus nebulosa). *Abstr., J. Parasit.* 20(2):136.
- _____. 1934. Some trematodes of the common bullhead (Ameiurus nebulosa). *Tr. Am. Micr. Soc.* 53(3):267-276.
- Baer, J. G. 1951. Ecology of animal parasites. Univ. of Illinois Press, Urbana, 224 p.
- Bailey, R. M. (chairman). 1960. A list of common and scientific names of fishes from the United States and Canada. *Am. Fish. Soc. Spec. Publ.* no. 2:102.
- Bangham, R. V. 1926. Parasites other than cestodes in black bass in Ohio. *Ohio J. Sci.* 25(6):255-270.
- _____. 1927. Diseases of fish in Ohio hatcheries. *Tr. Am. Fish. Soc.* 57:223-230.
- _____. 1928. Parasites of black bass. *Scient. Month.* 27(3):257-270.
- _____. 1937. Parasites of Wayne County (Ohio) stream fish. *Tr. Am. Fish. Soc.* 88:263-268.
- _____. 1941a. Parasites of fresh-water fish of southern Florida. *Proc. Florida Acad. Sci.* 5:289-307.
- _____. 1941b. Parasites from fish of Buckeye Lake, Ohio. *Ohio J. Sci.* 41(6):441-448.
- _____. 1941c. Parasites of fish of Algonquin Park Lakes. *Tr. Am. Fish. Soc.* 70:161-171.

- _____. 1951. Parasites of fish in the upper Snake River drainage and in Yellowstone Lake, Wyoming. *Zoologica, New York Zool. Soc.* 36(3):213-217.
- _____. 1955. Studies on fish parasites of Lake Huron and Manitoulin Island. *Am. Midl. Nat.* 53(1):184-194.
- _____, and G. W. Hunter. 1939. Studies on fish parasites of Lake Erie. Distribution studies. *Zoologica, New York Zool. Soc.* 24(4):385-448.
- _____, and C. E. Venard. 1942. Studies on parasites of Reel-foot Lake fish. IV. Distribution studies and checklist of parasites. *J. Tennessee Acad. Sci.* 17(1):22-38.
- Dogiel, V. A. 1958. Parasitology of fishes. Leningrad Univ. Press. (English transl. Z. Kabata. Oliver and Boyd, Edinburgh.)
- Essex, H. E., and G. W. Hunter, III. 1926. A biological study of fish parasites from the central states. *Tr. Ill. Acad. Sci.*
- Faust, E. C. 1925. Parasitism among the Helminths. *Am. Nat.* 59:197-519.
- Fischthal, J. H. 1947. Parasites of Northwest Wisconsin fishes. I. The 1944 survey. *Trans. Wisconsin Acad. Sci. Arts and Let.* 37:157-220.
- _____. 1950. Parasites of Northwest Wisconsin fishes. II. The 1945 survey. *Trans. Wisconsin Acad. Sci. Arts and Let.* 49(1):87-113.
- Hederlife, E. C. 1953. Parasites of the freshwater fishes of Northern California. Univ. of Calif. Press, Berkeley and Los Angeles, 110 p.
- Hoffman, G. L. 1960. Synopsis of strigeoidea (trematoda) of fishes and their life cycles. *BSFW, Fishery Bull.* 60:439-469.
- _____. 1963. Parasites of freshwater fish. I. Fungi. (*Saprolegnia* and relatives) of fish and fish eggs. *BSFW, Fishery Leaflet.* 564:1-6.
- _____. 1967. Parasites of North American Freshwater Fishes. Univ. of Calif. Press, Berkeley and Los Angeles, 308 p.
- Huggins, E. J. 1972. Parasites of fishes in South Dakota. *Bull.* 48L. Agric. Expt. Station, South Dakota State Univ.
- Hunter, G. W., III. 1927. Notes on the Caryophyllidae of North America. *J. Parasit.* 14(1):16-26.
- _____. 1932. An artificial lake with a low percentage of infected fish. *Tr. Am. Micr. Soc.* 51(1):22-27.
- _____. 1942. Studies on the parasites of freshwater fishes on Connecticut. *Conn. State Geol. and Nat. Hist. Survey. Bull. no. 63:* 228-288.

- _____, and W. S. Hunter. 1934. Further studies on fish and bird parasites. Suppl. 24th. Ann. Rept., New York State Conserv. Dept., No. IX, Rept. Biol. Survey Mohawk-Hudson Watershed. 267-283.
- Jenkins, R. M. 1957. The effect of gizzard shad on the fish populations of a small Oklahoma lake. Trans. Amer. Fish. Soc. 85:58-74.
- Jester, D. B., and B. L. Jensen. 1972. Life history and ecology of the gizzard shad (Dorosoma cepedianum), with special reference to Elephant Butte Lake. Agric. Expt. Station Research Rept. 218. Las Cruces, New Mexico.
- Kutkuhn, J. H. 1958. Utilization of gizzard shad by game fishes. Proc. Iowa Acad. Sci. 65:571-579.
- Leidy, J. 1856. A synopsis of Entozoa and some of their ectocongeners observed by the author. Proc. Acad. Nat. Sci. 8(1):42-58.
- Lewis, W. M. 1953. Analysis of the gizzard shad population of Crab Orchard Lake, Illinois. Tr. Ill. Acad. Sci. 46:231-234.
- Linton, E. 1897. Notes on Cestode parasites of fishes. Proc. U. S. Natl. Museum. 20:423-458.
- _____. 1898. Notes on Trematode parasites of fishes. Proc. U. S. Natl. Museum. 26:507-548.
- _____. 1914. On the seasonal distribution of fish parasites. Tr. Am. Fish. Soc. 44:48-56.
- McDaniel, J. S. 1963. Parasites from the genus Lepomis (Centrarchidae) in Lake Texoma, Oklahoma. Tr. Am. Micr. Soc. 82(4):423-425.
- Manter, H. W. 1926. Some North American fish Trematodes. Ill. Biol. Monog. 10(2):1-138.
- Marshall, W. S. and N. C. Gilbert. 1905. Notes on the food and parasites of some freshwater fishes from the lakes at Madison, Wisconsin. Rept. Comm. Fish. for 1905. 513-522.
- Meyer, F. P. 1958. Helminths of fishes from Trumbull Lake, Clay County, Iowa. Proc. Iowa Acad. Sci. 65:477-516.
- Miller, R. R. 1960. Systematics and Biology of the gizzard shad (Dorosoma cepedianum) and related fishes. Fish. Bull., Fish and Wildlife Service. 60:371-388.
- Muller, J. F. 1934. Parasites of Oneida Lake fishes. Ill. Biological and ecological survey of the worm parasites. IV. Additional notes on parasites of Oneida Lake fishes, including descriptions of new species. Roosevelt Wild Life Ann. 3(4):335-373.

- _____, and H. J. Van Cleave. 1932. Parasites of Oneida Lake fishes. II. Description of new species and some general taxonomic considerations especially concerning the trematode family Heterophyidae. *Roosevelt Wild Life Ann.* 3(2):79-137.
- Price, E. W. 1958. Some new monogenetic trematodes from the gizzard shad, (*Dorosoma cepedianum*). *J. Alabama Acad. Sci.* 30(2):68.
- _____. 1959. A proposed reclassification of the gastrocystoid monogenea. *J. Parasit.* 45(4):22-23.
- Putz, R. E., G. L. Hoffman, and C. E. Dunbar. 1965. Two new species of *Plistophora* (microsporidea) from North American fish with a synopsis of microsporidea of freshwater and Euryhaline fishes. *J. Protozool.* 12(2):228-236.
- Scheidermeyer, F. and W. M. Lewis. 1958. Utilization of gizzard shad by largemouth bass. *The Progressive Fish Culturist.* 18(3):137-138.
- Sroufe, S. A., Jr. 1958. *Mazocraeoides olentangiensis* n. sp., a monogenetic trematode parasitic on the gills of the gizzard shad (*Dorosoma cepedianum*). *J. Parasit.* 44(6):643-646.
- Van Cleave, H. J. 1916. Seasonal distribution of some Acanthocephala from fresh water hosts. *J. Parasit.* 2:106-110.
- _____. 1919. Acanthocephala from the Illinois River with descriptions of species and a synopsis of the family Neoechinorhynchidae. *Bull. Ill. Nat. Hist. Survey.* 131:222-257.
- _____. 1921. Acanthocephala from the eel. *Tr. Am. Micr. Soc.* 40:1-13.
- _____. 1923. Acanthocephala from the fishes of Oneida Lake, New York. *Roosevelt Wild Life Bull.* 2(1):73-86.
- _____. 1931. New Acanthocephala from fishes of Mississippi and a taxonomic reconsideration of forms with unusual numbers of cement glands. *Tr. Am. Micr. Soc.* 50:348-353.
- Ward, H. B. 1912. The distribution and frequency of animal parasites and parasitic diseases in North American freshwater fish. *Tr. Am. Fish. Soc.* 41:207-241.
- _____. 1918. Parasitic roundworms. In: Ward, H. B., and G. C. Whipple. *Freshwater Biology.* 506-552.
- _____, and T. B. Magath. 1916. Notes on some nematodes from freshwater fishes. *J. Parasit.* 3(2):57-64.
- Wilson, C. B. 1916. Copepod parasites of freshwater fishes and their economic relations to mussel glochidia. *Bull. U. S. Bur. Fish.* 34(1914):331-374.

- Yamaguti, S. 1959. *Systema Helminthum*. Vol. II. The Cestodes of vertebrates. Interscience, New York. 860 p.
- _____. 1961. *Systema Helminthum*. Vol. III. The Nematodes of vertebrates. Interscience, New York. 697 p.
- _____. 1963. *Systema Helminthum*. Vol. V. Acanthocephala. Interscience, New York. 421 p.
- _____. 1963. Parasitic copepoda and Branchiura of fishes. Interscience, New York. 1104 p.
- Yorke, W., and P. H. Maplestone: 1962. *The Nematode Parasites of Vertebrates*. Hafner Publ. Co., New York. 378 p.